

Generalized additive bases, König's lemma, and the Erdős-Turán conjecture*

Melvyn B. Nathanson[†]
 Department of Mathematics
 Lehman College (CUNY)
 Bronx, New York 10468
 Email: nathansn@alpha.lehman.cuny.edu

February 1, 2008

Abstract

Let A be a set of nonnegative integers. For every nonnegative integer n and positive integer h , let $r_A(n, h)$ denote the number of representations of n in the form $n = a_1 + a_2 + \cdots + a_h$, where $a_1, a_2, \dots, a_h \in A$ and $a_1 \leq a_2 \leq \cdots \leq a_h$. The infinite set A is called a *basis of order h* if $r_A(n, h) \geq 1$ for every nonnegative integer n . Erdős and Turán conjectured that $\limsup_{n \rightarrow \infty} r_A(n, 2) = \infty$ for every basis A of order 2. This paper introduces a new class of additive bases and a general additive problem, a special case of which is the Erdős-Turán conjecture. König's lemma on the existence of infinite paths in certain graphs is used to prove that this general problem is equivalent to a related problem about finite sets of nonnegative integers.

1 Representation functions and the Erdős-Turán conjecture

Let $\mathbf{N}_0$ and $\mathbf{Z}$ denote the nonnegative integers and integers, respectively. Let A be a finite set of integers. We denote the largest element of A by $\max(A)$ and the cardinality of A by $\text{card}(A)$. For any real numbers a and b , we denote by $[a, b]$ the finite set of integers n such that $a \leq n \leq b$.

For any set A of integers, we denote by $r_A(n, h)$ the number of representations of n in the form $n = a_1 + a_2 + \cdots + a_h$, where $a_1, a_2, \dots, a_h \in A$ and $a_1 \leq a_2 \leq \cdots \leq a_h$. The function r_A is called the *unordered representation function* of the set A , or, simply, the *representation function* of A .

*2000 Mathematics Subject Classification: 11B13, 11B34, 11B05. Key words and phrases. Additive bases, sumsets, representation functions, Erdős-Turán conjecture, König's lemma.

[†]This work was supported in part by grants from the NSA Mathematical Sciences Program and the PSC-CUNY Research Award Program.

The set A of nonnegative integers is called a *basis of order h* if every nonnegative integer can be represented as the sum of h not necessarily distinct elements of A . If A is a basis of order h with representation function r_A , then

$$1 \leq r_A(n, h) < \infty$$

for all nonnegative integers n . We call A an *asymptotic basis of order h* if $r_A(n, h) = 0$ for only finitely many $n \in \mathbf{N}_0$. If $h \geq 2$ and $f : \mathbf{N}_0 \rightarrow \mathbf{N}_0$ is any function such that $\text{card}(f^{-1}(0)) < \infty$, then the *representation function problem* is to determine if there exists a set A of nonnegative integers such that $r_A(n, h) = f(n)$ for all $n \geq 0$.

In the case of additive bases for the set of all integers, Nathanson [5] proved that every function is a representation function, that is, if $f : \mathbf{Z} \rightarrow \mathbf{N}_0 \cup \{\infty\}$ satisfies the condition $\text{card}(f^{-1}(0)) < \infty$, then for every $h \geq 2$ there exists a set A of integers such that $r_A(n, h) = f(n)$ for every integer n .

A special case of the representation function problem for nonnegative integers with $h = 2$ is the conjecture of Erdős and Turán [2] that the representation function $r_A(n, 2)$ of an asymptotic basis A of order 2 must be unbounded, that is,

$$\liminf_{n \rightarrow \infty} r_A(n, 2) > 0 \implies \limsup_{n \rightarrow \infty} r_A(n, 2) = \infty.$$

This is an important unsolved problem in additive number theory.

Dowd [1] and Grekos, Haddad, Helou, and Pikho [3] have given various equivalent formulations of the Erdős-Turán conjecture. In particular, Dowd proved that there exists a set A of nonnegative integers and a number c such that $r_A(n, 2) \in [1, c]$ for all nonnegative integers n if and only if for every N there exists a finite set A_N of nonnegative integers with $\max(A_N) \geq N$ and $r_A(n, 2) \in [1, c]$ for all $n = 0, 1, \dots, \max(A_N)$. In this paper we apply Dowd's method to obtain similar results for a new class of generalized additive bases.

2 Generalized additive bases

We extend the idea of an additive basis of order h as follows: Let $\mathcal{H} = \{H_n\}_{n=0}^{\infty}$ be a sequence of nonempty finite sets of positive integers. For any set A of nonnegative integers, we define the representation function

$$r_A(n, H_n) = \sum_{h_n \in H_n} r_A(n, h_n).$$

The set A of nonnegative integers will be called a *basis of order $\mathcal{H}$* if

$$r_A(n, H_n) \geq 1 \tag{1}$$

for all $n \geq 0$, and an *asymptotic basis of order $\mathcal{H}$* if the representation function satisfies (1) for all sufficiently large n .

Let $\mathcal{R} = \{R_n\}_{n=0}^{\infty}$ be a sequence of nonempty finite sets of positive integers. If

$$r_A(n, H_n) \in R_n \tag{2}$$

for every nonnegative integer n , then A will be called an $\mathcal{R}$ -basis of order $\mathcal{H}$. Since each R_n is a nonempty set of positive integers, it follows that every $\mathcal{R}$ -basis of order $\mathcal{H}$ is a basis of order $\mathcal{H}$. We shall call the set A an *asymptotic $\mathcal{R}$ -basis of order $\mathcal{H}$* if (2) holds for all sufficiently large n .

For any sequences $\mathcal{H} = \{H_n\}_{n=0}^\infty$ and $\mathcal{R} = \{R_n\}_{n=0}^\infty$ of nonempty finite sets of positive integers, we can ask if there exists an $\mathcal{R}$ -basis of order $\mathcal{H}$ or an asymptotic $\mathcal{R}$ -basis of order $\mathcal{H}$. This is the *generalized representation function problem*. The original Erdős-Turán conjecture corresponds to the special case $H_n = \{2\}$ and $R_n = [1, c]$ for all $n \geq 0$. It is an open problem to determine the number of distinct $\mathcal{R}$ -bases of order $\mathcal{H}$ and asymptotic $\mathcal{R}$ -bases of order $\mathcal{H}$ for a given pair of sequences $\mathcal{H}$ and $\mathcal{R}$.

Let $h \geq 2$ and let f be a function such that $f(n)$ is a positive integer for every nonnegative integer n . We introduce the sets $H_n = \{h\}$ and $R_n = \{f(n)\}$ for all n , and the sequences $\mathcal{H} = \{H_n\}_{n=0}^\infty$ and $\mathcal{R} = \{R_n\}_{n=0}^\infty$. Then an $\mathcal{R}$ -basis of order $\mathcal{H}$ is a basis A of order h whose representation function satisfies $r_A(n, h) = f(n)$ for all $n \in \mathbf{N}_0$, and so the representation function problem for bases of order h is a special case of the generalized representation function problem.

An $\mathcal{R}$ -basis of order $\mathcal{H}$ is not necessarily infinite. For example, if $H_0 = R_0 = \{1\}$ and if $H_n = \{n\}$ and $1 \in R_n$ for all $n \geq 1$, then the set $\{0, 1\}$ is an $\mathcal{R}$ -basis of order $\mathcal{H}$.

Theorem 1 *Let $\mathcal{H} = \{H_n\}_{n=0}^\infty$ be a sequence of nonempty finite sets of positive integers. There exists a finite set A that is a basis of order $\mathcal{H}$ or an asymptotic basis of order $\mathcal{H}$ if and only if*

$$\liminf_{n \rightarrow \infty} \frac{\max(H_n)}{n} > 0.$$

Proof. Let $h_n^* = \max(H_n)$. Let A be a finite set of nonnegative integers that is a basis of order $\mathcal{H}$. Then $0, 1 \in A$ and so $\max(A) \geq 1$. Every positive integer n can be represented as the sum of h_n elements of A for some $h_n \in H_n$, and so

$$n \leq h_n \max(A) \leq h_n^* \max(A).$$

It follows that

$$\liminf_{n \rightarrow \infty} \frac{h_n^*}{n} \geq \frac{1}{\max(A)} > 0.$$

Conversely, if $\liminf_{n \rightarrow \infty} h_n^*/n > 0$, then there exists a positive integer m such that

$$h_n^* \geq \frac{n}{m}$$

for all $n \geq 0$. Consider the finite set $A = [0, m]$. By the division algorithm, every positive integer n can be written in the form $n = qm + r$, where q and r are nonnegative integers and $0 \leq r \leq m - 1$. If $r = 0$, then $q = n/m \leq h_n^*$ and

$$n = q \cdot m + (h_n^* - q) \cdot 0 \in h_n^* A.$$

If $1 \leq r \leq m-1$, then $q = (n-r)/m < h_n^*$. Since h_n^* and q are integers, it follows that $h_n^* \geq q+1$ and

$$n = q \cdot m + 1 \cdot r + (h_n^* - q - 1) \cdot 0 \in h_n^* A.$$

In both cases, $r_A(n, H_n) \geq r_A(n, h_n^*) \geq 1$, and the finite set A is a basis of order $\mathcal{H}$.

If A is an asymptotic basis of order $\mathcal{H}$, then $r_A(n, H_n) = 0$ for only finitely many $n \in \mathbf{N}_0$, and so there is a finite set F of nonnegative integers such that $A \cup F$ is a basis of order $\mathcal{H}$. Therefore, there exists a finite set that is a basis of order $\mathcal{H}$ if and only if there exists a finite set that is an asymptotic basis of order $\mathcal{H}$. This completes the proof. $\square$

Let $\mathcal{H} = \{H_n\}_{n=0}^\infty$ and $\mathcal{R} = \{R_n\}_{n=0}^\infty$ be sequences of nonempty finite sets of positive integers. A nonempty finite set A of nonnegative integers will be called a *finite basis of order $\mathcal{H}$* if

$$r_A(n, H_n) \geq 1$$

for all $n \in [0, \max(A)]$, and a *finite $\mathcal{R}$ -basis of order $\mathcal{H}$* if

$$r_A(n, H_n) \in R_n$$

for all $n \in [0, \max(A)]$.

Theorem 2 *Let $\mathcal{H} = \{H_n\}_{n=0}^\infty$ and $\mathcal{R} = \{R_n\}_{n=0}^\infty$ be sequences of nonempty finite sets of positive integers.*

- (i) *If A is a basis of order $\mathcal{H}$, or if A is a finite basis of order $\mathcal{H}$ with $\max(A) \geq 1$, then $0, 1 \in A$.*
- (ii) *If A is an $\mathcal{R}$ -basis of order $\mathcal{H}$ or if A is a finite $\mathcal{R}$ -basis of order $\mathcal{H}$ with $\max(A) \geq 1$, then $\text{card}(H_0) \in R_0$ and $\text{card}(H_1) \in R_1$.*
- (iii) *If A is an $\mathcal{R}$ -basis of order $\mathcal{H}$, then $A_N = A \cap [0, N]$ is a finite $\mathcal{R}$ -basis of order $\mathcal{H}$ for every $N \geq 0$.*
- (iv) *If $A \neq \{0\}$ is a finite $\mathcal{R}$ -basis of order $\mathcal{H}$, then $F' = F \setminus \{\max(F)\}$ is also a finite $\mathcal{R}$ -basis of order $\mathcal{H}$.*

Proof. To prove (i) and (ii), we observe that if $r_A(0, H_0) \geq 1$, then $0 \in A$. If $r_A(1, H_1) \geq 1$, then $1 \in A$. Since, for every $h \geq 1$, both 0 and 1 have unique representations as sums of exactly h nonnegative integers, it follows that if $r_A(0, H_0) \in R_0$, then

$$r_A(0, H_0) = \sum_{h_0 \in H_0} r_A(0, h_0) = \sum_{h_0 \in H_0} 1 = \text{card}(H_0) \in R_0.$$

Similarly, if $r_A(1, H_1) \in R_1$, then

$$r_A(1, H_1) = \text{card}(H_1) \in R_1.$$

The statements (iii) and (iv) follow immediately from the definition of a finite basis. $\square$

3 König's lemma

The principal tool in this paper is König's lemma on the existence of infinite paths in trees. For completeness, we include a short proof below.

A *graph* G consists of a nonempty set $\{v\}$, whose elements are called *vertices*, and a set $\{e\}$, whose elements are called *edges*. Each edge is a set $e = \{v, v'\}$, where v and v' are vertices and $v \neq v'$. Thus, we are considering only graphs without loops or multiple edges.

We use the following terminology. The vertices v and v' are called *adjacent* if $\{v, v'\}$ is an edge. The degree of a vertex v is the number of edges e with $v \in e$. A *path* in G from vertex v to vertex v' is a sequence of vertices $v_0, v_1, v_2, \dots, v_n$ such that $v_0 = v$, $v_n = v'$, and v_{i-1} is adjacent to v_i for all $i = 1, \dots, n$. We define the *length* of this path by n . The graph G is *connected* if for every two vertices v and v' with $v \neq v'$ there is a path from v to v' . A graph is connected if and only if, for some vertex v_0 , there is a path from v_0 to v for every vertex $v \neq v_0$.

A *simple path* in G is a path whose vertices are pairwise distinct. A *simple circuit* is a sequence of vertices $v_0, v_1, v_2, \dots, v_n$ such that $n \geq 3$, $\{v_{i-1}, v_i\}$ is an edge for $i = 1, \dots, n$, $v_i \neq v_j$ for $0 \leq i < j \leq n-1$, and $v_0 = v_n$. A graph G has no simple circuits if and only, for every pair of distinct vertices v and v' , there is at most one simple path from v to v' . An *infinite simple path* is an infinite sequence of pairwise distinct vertices $v_0, v_1, v_2, \dots$ such that v_{i-1} is adjacent to v_i for all $i \geq 1$.

A *tree* is a connected graph with no simple circuits. A *rooted tree* is a tree with a distinguished vertex, called the *root* of the tree. In a rooted tree, for every vertex v different from the root, there is a unique simple path in the tree from the root to v .

Theorem 3 (König's lemma) *If T is a rooted tree with infinitely many vertices such that every vertex has finite degree, then T contains an infinite simple path beginning at the root.*

Proof. Let v_0 be the root of the tree. We use induction to prove that for every n there is a simple path $v_0, v_1, \dots, v_n$ such that the tree T contains infinitely many vertices v for which the unique simple path from the root v_0 to v begins with the vertices $v_0, v_1, \dots, v_n$. Since T has infinitely many vertices, the root v_0 satisfies this condition.

Let $n \geq 1$, and assume that we have constructed a simple path $v_0, v_1, \dots, v_{n-1}$ of vertices of the tree T with the property that T contains an infinite set I_{n-1} of vertices such that, for every $v \in I_{n-1}$, the unique simple path from v_0 to v passes through vertex v_{n-1} . Since the degree of v_{n-1} is finite, the set F_n of vertices $v \neq v_{n-1}$ that are adjacent to v_{n-1} is a finite set. For every vertex $v \in I_{n-1}$, there is a unique simple path in T that begins at v_0 , passes through v_{n-1} and exactly one of the vertices in F_n , and ends at v . By the pigeonhole principle, since I_{n-1} is infinite, there is a vertex $v_n \in F_n$ and an infinite set $I_n \subseteq I_{n-1}$ of vertices such that, for every $v \in I_n$, the unique path from v_0 to v

passes through v_n . This completes the induction. The vertices $v_0, v_1, v_2, \dots$ are pairwise distinct, and $v_0, v_1, v_2, \dots$ is an infinite simple path in T . $\square$

4 The generalized representation function problem

In this section we prove that there exists an infinite $\mathcal{R}$ -basis of order $\mathcal{H}$ if and only if there exist arbitrarily large finite $\mathcal{R}$ -bases of order $\mathcal{H}$.

Theorem 4 *Let $\mathcal{R} = \{R_n\}_{n=0}^\infty$ and $\mathcal{H} = \{H_n\}_{n=0}^\infty$ be sequences of nonempty finite sets of positive integers such that*

$$\lim_{n \rightarrow \infty} \frac{\max(H_n)}{n} = 0. \quad (3)$$

There exists an $\mathcal{R}$ -basis A of order $\mathcal{H}$ if and only if for every N there exists a finite $\mathcal{R}$ -basis A_N of order $\mathcal{H}$ with $\max(A_N) \geq N$.

Proof. If A is a $\mathcal{R}$ -basis of order $\mathcal{H}$, then, by (3) and Theorem 1, the set A is infinite, hence for every N there is an integer $a(N) \in A$ with $a(N) \geq N$. By Theorem 2, the set $A_N = A \cap [0, a(N)]$ is a finite $\mathcal{R}$ -basis of order $\mathcal{H}$ with $\max(A_N) \geq N$.

Conversely, suppose that for every N there exists a finite $\mathcal{R}$ -basis A_N of order $\mathcal{H}$ with $\max(A_N) \geq N$. If $N \geq 1$, then $0, 1 \in A_N$ and the sets $\{0\}$ and $\{0, 1\}$ are finite $\mathcal{R}$ -bases of order $\mathcal{H}$.

We construct the graph T whose vertices are the finite $\mathcal{R}$ -bases of order $\mathcal{H}$. This graph has infinitely many vertices, since there are finite $\mathcal{R}$ -bases of order $\mathcal{H}$ with arbitrarily large maximum elements.

Vertices V and V' will be called adjacent in this graph if $V' \subseteq V$ and $V \setminus V' = \{\max(V)\}$. The sets $\{0\}$ and $\{0, 1\}$ are adjacent vertices of this graph, and $\{0, 1\}$ is the only vertex adjacent to $\{0\}$. If V is a vertex and $\text{card}(V) \geq 2$, then it follows from Theorem 2 that $V' = V \setminus \{\max(V)\}$ is a vertex. Moreover, V' is the unique vertex adjacent to V in T such that $\text{card}(V') = \text{card}(V) - 1$. If V'' is adjacent to V and $V'' \neq V'$, then $V = V'' \setminus \{\max(V'')\}$ and $\text{card}(V'') = \text{card}(V) + 1$.

We shall prove that T is a rooted tree with root $V_0 = \{0\}$. Let $V = \{a_0, a_1, \dots, a_n\}$ be a vertex, where $0 = a_0 < a_1 < \dots < a_n$. For every $k = 0, 1, \dots, n$, the set $V_k = \{a_0, a_1, \dots, a_k\}$ is a finite $\mathcal{R}$ -basis of order $\mathcal{H}$, hence is a vertex of T . Then $V_0 = \{0\}$, $V_n = V$, and $V_0, V_1, \dots, V_n$ is a simple path in T from the root V_0 to V . It follows that the graph T is connected.

Suppose that $n \geq 3$ and $V_0, V_1, \dots, V_{n-1}, V_n$ is a simple circuit in T , where $V_n = V_0$. Let $V_{n+1} = V_1$. Since each vertex is a finite set of integers, we can choose $k \in [1, n]$ such that V_k is a vertex in the circuit of maximum cardinality. Vertices V_{k-1} and V_{k+1} are adjacent to V_k , hence $\text{card}(V_k) - \text{card}(V_{k-1}) = \pm 1$ and $\text{card}(V_k) - \text{card}(V_{k+1}) = \pm 1$. The maximality of $\text{card}(V_k)$ implies

that $\text{card}(V_k) - \text{card}(V_{k-1}) = \text{card}(V_k) - \text{card}(V_{k+1}) = 1$, and so $V_{k-1} = V_k \setminus \{\max(V_k)\} = V_{k+1}$, which is impossible. Therefore, T contains no simple circuit, and so T is a tree.

To apply König's lemma, we must prove that every vertex of this tree has finite degree. The only vertex adjacent to the root $\{0\}$ is $\{0, 1\}$, hence $\{0\}$ has finite degree. Let $V \neq \{0\}$ be a vertex of T . Then $1 \in V$ and so $\max(V) \geq 1$. Suppose that the V is adjacent to infinitely many vertices V' . The only subset of V that is a vertex adjacent to V is $V \setminus \{\max(V)\}$. Every other vertex V' adjacent to V is a superset of V of the form $V' = V \cup \{\max(V')\}$. For each such V' , the integer $n = \max(V') - 1$ must be an element of the sumset $h_n V$ for some $h_n \in H_n$, and so

$$n \leq h_n \max(V) \leq \max(H_n) \max(V).$$

Since $n \in h_n V$ for infinitely many integers n , it follows that

$$\limsup_{n \rightarrow \infty} \frac{\max(H_n)}{n} \geq \frac{1}{\max(V)} > 0,$$

which contradicts (3). Thus, every vertex of the infinite tree T has finite degree. By König's lemma, the tree must contain an infinite simple path $\{0\} = V_0, V_1, V_2, \dots$. For each nonnegative integer n , let $a_n = \max(V_n)$. Then $V_n = \{a_0, a_1, \dots, a_n\}$ for all $n = 0, 1, 2, \dots$. Let

$$A = \{a_n\}_{n=0}^{\infty} = \bigcup_{n=0}^{\infty} V_n.$$

Since $a_n \geq n$, it follows that

$$r_A(n, H_n) = \sum_{h_n \in H_n} r_A(n, h_n) = \sum_{h_n \in H_n} r_{V_n}(n, h_n) = r_{V_n}(n, H_n) \in R_n,$$

and so A is a $\mathcal{R}$ -basis of order $\mathcal{H}$. This completes the proof. $\square$

Theorem 5 *Let $h \geq 2$ and let f be a function such that $f(n)$ is a positive integer for every nonnegative integer n . There exists a basis A of order h with representation function $r_A(n, h) = f(n)$ if and only if for every N there exists a finite set A_N of nonnegative integers with $\max(A_N) \geq N$ and $r_A(n, h) = f(n)$ for all $n = 0, 1, \dots, \max(A_N)$.*

Proof. This follows immediately from Theorem 4 with $H_n = \{h\}$ and $R_n = \{f(n)\}$ for all nonnegative integers n . $\square$

Applying Theorem 4 to the classical Erdős-Turán conjecture, we obtain the following result of Dowd [1, Theorem 2.1].

Theorem 6 *Let $c \geq 1$ and $h \geq 2$. There exists a basis A of order h such that*

$$r_A(n, h) \leq c \quad \text{for all } n \geq 0$$

if and only if, for every N , there exists a finite set A_N of nonnegative integers such that $\max(A_N) \geq N$ and

$$1 \leq r_{A_N}(n, h) \leq c$$

for all $n = 0, 1, \dots, \max(A_N)$.

Proof. This follows immediately from Theorem 4 with $H_n = \{h\}$ and $R_n = [1, c]$ for all $n \geq 0$. $\square$

5 Ordered representation functions

There are other important representation functions in additive number theory. For example, for any set A of integers, the *ordered representation function* $r'_A(n, h)$ counts the number of h -tuples $(a_1, \dots, a_h) \in A^h$ such that $a_1 + \dots + a_h = n$. Nathanson [4] proved the following uniqueness theorem for ordered representation functions: For any function $f : \mathbf{N}_0 \rightarrow \mathbf{N}_0$ and for any positive integer h , there exists at most one set A of nonnegative integers such that $r'_A(n, h) = f(n)$ for all $n \in \mathbf{N}_0$. He also showed that uniqueness does not hold if ordered representation functions only eventually coincide, and he described all pairs of sets A and B of nonnegative integers such that $r'_A(n, 2) = r'_B(n, 2)$ for all sufficiently large integers n .

We can define *basis of order $\mathcal{H}$* and *$\mathcal{R}$ -basis of order $\mathcal{H}$* in terms of the ordered representation function. Theorem 4 is also true for ordered representation functions.

References

- [1] M. Dowd, *Questions related to the Erdős-Turán conjecture*, SIAM J. Discrete Math. **1** (1988), 142–150.
- [2] P. Erdős and P. Turán, *On a problem of Sidon in additive number theory and some related questions*, J. London Math. Soc. **16** (1941), 212–215.
- [3] G. Grekos, L. Haddad, C. Helou, and J. Pihko, *On the Erdős-Turán conjecture*, Preprint, 2002.
- [4] M. B. Nathanson, *Representation functions of sequences in additive number theory*, Proc. Amer. Math. Soc. **72** (1978), 16–20.
- [5] M. B. Nathanson, *Every function is the representation function of an additive basis for the integers*, Preprint, 2003.